

Recent Update on MODIS/VIIRS Deep Blue Data Continuity and New Aerosol Products




*Photo taken from Space Shuttle:
Fierce dust front over Libya*

*N. Christina Hsu (PI), Andrew M. Sayer, Jaehwa Lee,
Corey Bettenhausen, N. Carletta, and Si-Chee Tsay*

Laboratory for Atmospheres

NASA Goddard Space Flight Center, Greenbelt, Maryland USA




Recent Progress on Deep Blue Aerosol Algorithm for VIIRS


- ***Expand coverage from **arid and semi-arid** regions into **vegetated** (SeaWiFS, MODIS C6, and VIIRS) areas as well as **oceans** (SeaWiFS and VIIRS only)***
- ***Develop and employ **non-spherical dust models** for aerosol retrievals***
- ***Improve cloud screening to distinguish **heavy smoke plumes** from clouds***
- ***Account for the effect of **water vapor** in identifying **strongly absorbing mineral dust** using the **IR channels*****

VIIRS 2012-2016 Seasonal 550nm AOD Maps


Winter




Spring



Summer




Fall



Aerosol Optical Depth




VIIRS ocean retrieval algorithm



- The VIIRS ocean algorithm is an extension and improvement on our SeaWiFS algorithm
 - Similar in principle to other common approaches (e.g. MODIS) as well
- Retrieve AOD, fine mode fraction (Ångström exponent), aerosol type (from a selection of models)
 - Includes nonspherical dust model
- Cloud screening seems effective even in cases of heavy aerosol loading

Effects of VIIRS Vicarious Calibrations on AOD retrievals







- Applying these corrections removes the bulk of the AOD bias over water
- Suggest MODIS absolute calibration is closer to the truth than VIIRS

For more details, see our poster (Atmospheres section): Sayer *et al.*, “Vicarious calibration of S-NPP VIIRS reflective solar M-bands against MODIS Aqua over dark water scenes”




Comparisons of VIIRS ocean products with Maritime Aerosol Network (MAN)

- AOD retrieval quality similar to, or better than, standard MODIS product
- Ångström exponent (AE) and fine mode AOD fraction compare favorably to MAN data, even when the AOD is not high
- Colors indicate aerosol optical model: retrieved, not prescribed
 - Either **marine**, **dust**, or **fine-dominated**






VIIRS Comparisons with AERONET AOT over **Land**




**Urban
Pollution**



**Urban
Pollution,
Dust,
Smoke**







Sahel



Long-Term Data Continuity From MODIS to VIIRS

Comparisons of Daily VIIRS vs. MODIS Regional AOD





- The AOD values from VIIRS and MODIS are comparable on regional scales; An important step toward a consistent aerosol data record from MODIS to VIIRS

Planning for MODIS Collection 7:


- (1) Provide new *Aerosol Plume Height* product using combined MODIS and OMI data
- (2) Extend *Deep Blue Aerosol* products from Cloud free to Cloudy regions

Validation using data from 7SEAS field experiment over Southeast Asia


For more details, see our poster: Lee *et al.*, Retrieving the height of smoke and dust aerosols by synergistic use of multiple satellite sensors




CALIOP Total Attenuated Backscatter



Doi_Ang_Khang - 29 MAR 2013



Doi_Ang_Khang - 29 MAR 2013




Planning for MODIS Collection 7:


- (1) Provide new *Aerosol Plume Height* product using combined MODIS and OMI data
- (2) Extend *Deep Blue* Aerosol products from Cloud free to Cloudy regions

New Deep Blue Aerosol Products for MODIS C7: AOD and Aerosol Forcing above Clouds


Aqua/MODIS RGB
March 6, 2009





MODIS C6 Deep Blue AOD



MODIS C7 Deep Blue
new AOD above clouds




CERES TOA SW Flux (Wm^{-2}) **MODIS Deep Blue Aerosol Forcing (Wm^{-2})**




100 240 380 520 660 800

0 40 80 120 160 200

(a) MODIS Terra 13 Sep 2000 09:25




(b) AATS profile




- Validation case studies: take advantage of field campaign observations of aerosols above clouds
 - Observations from the Ames Airborne Tracking Sunphotometer (AATS; <http://geo.arc.nasa.gov/sgg/AATS-website/>)
 - For our purposes, a bit like an AERONET Sun photometer on a plane
 - Example from SAFARI-2000: smoke over a flat (~1 km) stratocumulus deck off the coast of Namibia


(a) MODIS Terra 13 Sep 2000 09:25




(b) C6 COD (liquid)



(c) AAC AOD



(d) AAC COD



- AATS 550 nm AOD: **0.49+/-0.02**; our MODIS AOD retrieval: **0.51+/-0.09**
- Our COD is well-correlated with but ~50% higher than operational C6 product, consistent with strong absorption by the smoke

Reference: Sayer et al., 2016, JGR, Extending 'Deep Blue' aerosol retrieval coverage to cases of absorbing aerosols above clouds: sensitivity analysis and first case studies

➤ Selected as a Research Spotlight by AGU: <https://eos.org/research-spotlights/measuring-atmospheric-aerosols-despite-the-clouds>

Extending Deep Blue Aerosol Products From LEO to GEO: Preliminary Results Using Himawari-8 Observations



Summary

- Based upon the comparisons with AERONET AOD global observations, the expected error for VIIRS DB is $0.05 \pm 20\%$ over land and $0.03 \pm 10\%$ over ocean, which is comparable to that for MODIS DB. The AOD time series from VIIRS and MODIS are consistent with each other.
- Thanks for the support from Atmosphere SIPS at Wisconsin, VIIRS version 1 DB aerosol products will be ready for release to the public in the summer of 2016.
- We have started planning for the MODIS C7 reprocessing to implement aerosol plume height and aerosol above cloud retrievals into the Deep Blue algorithm. The corresponding algorithm descriptions and preliminary validation results have been published in peer-reviewed journals.